

Show this card to any healthcare professional involved in your medical care – for example, your dentist or an emergency doctor.

Read the patient information leaflet for more information or scan the QR-code below.



www.fimea.fi

Finnish Medicines Agency Fimea
Register for adverse drug reactions
P.O. Box 55, FI-00034 FIMEA

Your name:

Name of physician who prescribed **RINVOQ**[®]:

Physician's phone number:

The date you started **RINVOQ**[®]:

Version 2.0. Date approved June, 2021.

FI-RNQ-210029/RMP/5.10.2021

AbbVie Oy
Tel. 010 2411 200

abbvie

PATIENT ALERT CARD

Keep this card with you all the time

Safety Information
about **RINVOQ**[®]
(upadacitinib) for Patients



This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effect you may get to Fimea: www.fimea.fi or to the local representative of the marketing authorisation holder (AbbVie Oy).

This card contains important safety information you should be aware of – before and during treatment with **RINVOQ**[®].

Risk of infections

RINVOQ® may make an existing infection worse or increase the chance of you getting a new infection – for example tuberculosis (TB), shingles or viral hepatitis.

Tell your doctor straight away if you notice signs of infection, such as:

- Fever, sweating, chills, weight loss, or a cough that won't go away – these may be signs of TB
- Painful skin rash with blisters – this may be a sign of shingles
- Feeling tired or short of breath – this may be a sign of pneumonia.

Vaccines – used to help prevent infections

Live vaccines (for example influenza vaccine by nasal spray, varicella, measles/mumps/rubella) should not be given during RINVOQ® treatment, or just before starting RINVOQ® treatment.

Before being given any vaccines, talk to your doctor – they will know which vaccines you should **not** be given before or during treatment with RINVOQ®.

Cholesterol

Your doctor will check your cholesterol levels while you are taking RINVOQ® – this is to decide if you need to start a medicine to lower your cholesterol levels.

Contraception, pregnancy, and breast-feeding

RINVOQ® must not be taken during pregnancy.

- Use effective contraception while taking RINVOQ® – and for 4 weeks after your last dose. Talk to your doctor about effective contraception.
- Tell your doctor straight away if you wish to become pregnant, or if you become pregnant.
- Do not breast-feed while using RINVOQ®.

Blood clots in veins (DVT) or lungs (PE)

Tell your doctor straight away if you get signs of DVT or PE, such as a painful swollen leg or chest pain.