

## **Part VI: Summary of the risk management plan**

### **Summary of risk management plan for Sitagliptin/Metformin STADA 50 mg/850 mg film-coated tablets and 50 mg/1 000 mg film-coated tablets (Sitagliptin/Metformin)**

This is a summary of the risk management plan (RMP) for Sitagliptin/Metformin STADA. The RMP details important risks of Sitagliptin/Metformin STADA, how these risks can be minimised, and how more information will be obtained about Sitagliptin/Metformin STADA's risks and uncertainties (missing information).

Sitagliptin/Metformin STADA's summary of product characteristics (SmPC) and its package leaflet give essential information to healthcare professionals and patients on how Sitagliptin/Metformin STADA should be used.

#### **I. The medicine and what it is used for**

Sitagliptin/Metformin STADA is authorised for adult patients with type 2 diabetes mellitus:

Sitagliptin/Metformin STADA is indicated as an adjunct to diet and exercise to improve glycaemic control in patients inadequately controlled on their maximal tolerated dose of metformin alone or those already being treated with the combination of sitagliptin and metformin.

Sitagliptin/Metformin STADA is indicated in combination with a sulphonylurea (i.e., triple combination therapy) as an adjunct to diet and exercise in patients inadequately controlled on their maximal tolerated dose of metformin and a sulphonylurea.

Sitagliptin/Metformin STADA is indicated as triple combination therapy with a peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) agonist (i.e., a thiazolidinedione) as an adjunct to diet and exercise in patients inadequately controlled on their maximal tolerated dose of metformin and a PPAR $\gamma$  agonist.

Sitagliptin/Metformin STADA is also indicated as add-on to insulin (i.e., triple combination therapy) as an adjunct to diet and exercise to improve glycaemic control in patients when stable dose of insulin and metformin alone do not provide adequate glycaemic control (see SmPC for the full indication). It contains sitagliptin and metformin as the active substances and it is given orally.

#### **II. Risks associated with the medicine and activities to minimise or further characterise the risks**

Important risks of Sitagliptin/Metformin STADA, together with measures to minimise such risks and the proposed studies for learning more about Sitagliptin/Metformin STADA's risks, are outlined below.

Measures to minimise the risks identified for medicinal products can be:

- Specific information, such as warnings, precautions, and advice on correct use, in the package leaflet and SmPC addressed to patients and healthcare professionals;

- Important advice on the medicine’s packaging;
- The authorised pack size — the amount of medicine in a pack is chosen so to ensure that the medicine is used correctly;
- The medicine’s legal status — the way a medicine is supplied to the patient (e.g. with or without prescription) can help to minimise its risks.

Together, these measures constitute *routine risk minimisation* measures.

If important information that may affect the safe use of Sitagliptin/Metformin STADA is not yet available, it is listed under ‘missing information’ below.

### ***II.A List of important risks and missing information***

Important risks of Sitagliptin/Metformin STADA are risks that need special risk management activities to further investigate or minimise the risk, so that the medicinal product can be safely taken. Important risks can be regarded as identified or potential. Identified risks are concerns for which there is sufficient proof of a link with the use of Sitagliptin/Metformin STADA. Potential risks are concerns for which an association with the use of this medicine is possible based on available data, but this association has not been established yet and needs further evaluation. Missing information refers to information on the safety of the medicinal product that is currently missing and needs to be collected (e.g. on the long-term use of the medicine);

| <b>Summary of safety concerns</b> |   |
|-----------------------------------|---|
| Important identified risks        | <ul style="list-style-type: none"> <li>• Lactic acidosis</li> </ul>                         |
| Important potential risks         | <ul style="list-style-type: none"> <li>• Pancreatic cancer</li> </ul>                       |
| Missing information               | <ul style="list-style-type: none"> <li>• Exposure during pregnancy and lactation</li> </ul> |

### ***II.B Summary of important risks***

The safety information in the proposed Product Information is aligned to the reference medicinal product.

### ***II.C Post-authorisation development plan***

#### **II.C.1 Studies which are conditions of the marketing authorisation**

There are no studies which are conditions of the marketing authorisation or specific obligation of Sitagliptin/Metformin STADA.

#### **II.C.2 Other studies in post-authorisation development plan**

There are no studies required for Sitagliptin/Metformin STADA.